



LABORATOIRE DE PROBABILITÉS
STATISTIQUE & MODÉLISATION



Identifiability of VAR(1) model in a stationary setting

Bixuan LIU

Supervised by: Stéphane Robin, Viet Chi Tran and Elisa Thébault

Algebraic statistics

Lille, March 12th-13th, 2026

Overview

1. Introduction
2. Identifiability framework
 - Statistical model
 - Identifiability definition
 - Identifiability criteria
3. Main results
4. Ongoing work

Introduction

Ecological background

Ecosystems are often modeled by **dynamical systems**.

For example, Robert May's model¹

$$\frac{d\mathbf{x}_t}{dt} = \Lambda^T \mathbf{x}_t,$$

- $\mathbf{x}_t$: n -dimensional vector of species abundances
- Λ : **deterministic** $n \times n$ interaction matrix
 - cooperation, competition, predation

(and the Lotka-Volterra equations²)

Central topic in ecology: determination of Λ .

1. May (1972) 2. Akjouj et al. (2024)

Ecological background

Central topic in ecology: determination of Λ .

Sampling insufficiency problem¹

Time-series sampling is **too costly** and are often sacrificed in practice. Instead, the limited resources are used to cover as much ground as possible.

Example: Study of a global ocean cross-domain plankton co-occurrence network with one-time-samplings of 200 stations².

Subject of this work

Recovery of dynamic networks in ecosystems using only **static data**^(*).

(*) e.g. i.i.d. samples of abundances of the same network from multiple sites.

1. Legendre et al. (2010) 2. Chaffron et al. (2021)

VAR(1)

The First-order Vector Autoregressive (VAR(1)) model:

$$\begin{cases} \mathbf{x}_1 = \boldsymbol{\epsilon}_1, \\ \mathbf{x}_t = \Lambda^T \mathbf{x}_{t-1} + \boldsymbol{\epsilon}_t \quad \text{for } t > 1, \\ \boldsymbol{\epsilon}_t \sim \mathcal{N}(0, \omega I_n) \end{cases}$$


where Λ is a **deterministic** $n \times n$ matrix.

Corresponding graph


$G = (V, \mathfrak{E}_G)$, where

- $V = \{1, \dots, n\}$, representing n interacting random variables,
- $\mathfrak{E}_G$ contains ordered pairs (i, j) s.t. $\Lambda_{ij} \neq 0$.

Graphical model



(a) The graphical VAR(1) model



(b) The encoded graph G

VAR(1) in a stationary setting

VAR(1) model:

$$\begin{cases} \mathbf{x}_1 = \boldsymbol{\epsilon}_1, \\ \mathbf{x}_t = \Lambda^T \mathbf{x}_{t-1} + \boldsymbol{\epsilon}_t \quad \text{for } t > 1, \\ \boldsymbol{\epsilon}_t \sim \mathcal{N}(0, \omega I_n) \end{cases}$$

If $\rho(\Lambda) < 1$, then $\mathbf{x}_t$ admits a stationary distribution $\mathbf{x}_\infty \sim \mathcal{N}(0, \Sigma)$ where

$$\Sigma = \Lambda^T \Sigma \Lambda + \omega I_n.$$

Assume Σ is known, can we recover the support of Λ , i.e. the underlying graph G ?

Obstacles

1. **Lack of reference:** VAR(1) graph recovery in the stationary regime is **rarely studied**; most work assumes time-series data^{1,2}.
2. **Classical methods not applicable:** The VAR(1) interaction graph **differs** from the graphical model of its stationary distribution. $\Rightarrow$ Results from Gaussian Graphical Models are **irrelevant**.

1. Gong et al. (2015) 2. Tank et al. (2019)

Related work

Recall

$$\Sigma = \Lambda^T \Sigma \Lambda + \omega I_n.$$

Young's condition¹ - counting the parameters

A necessary condition for identifiability is that the number of non-zero off-diagonal elements of Λ be no more than $n(n-3)/2$ and that $n \geq 5$.

Young's condition is too weak and has no guarantee for identifiability.

Drton's work²

Algebraic approaches are applied to linear structural equation models (SEMs), and sufficient graphical conditions for identifiability are derived.

1. Young et al. (2019) 2. Drton (2025)

Identifiability framework

Statistical model

The parametrization map:

$$\phi_G : (\Lambda, \omega) \mapsto \Sigma, \text{ s.t. } \Sigma = \Lambda^T \Sigma \Lambda + \omega I_n$$

is a well defined, fixed-point type function.

The stationary VAR(1) model

Given a directed graph G , and

$$M_G = \{\Lambda \mid \rho(\Lambda) < 1, \text{ and } \Lambda_{ij} = 0 \text{ if } (i, j) \notin \mathfrak{E}_G\},$$

then the stationary VAR(1) model is defined as

$$\mathbf{M}_G = \{\Sigma \mid \Sigma = \Lambda^T \Sigma \Lambda + \omega I_n, \Lambda \in M_G, \omega \in \mathbb{R}^+\}.$$

- $\mathbf{M}_G$ is (i) the image of ϕ_G and (ii) the set of covariance matrices Σ that can be associated with a matrix Λ of support G .

Global identifiability

Identifiability of a **finite family** $\Rightarrow$ identifiability of pairs.

Global identifiability

$$\mathbf{M}_1 \cap \mathbf{M}_2 = \emptyset.$$

$\Rightarrow$ too **strong**. Example: fix $\omega = 1$,

$$\Lambda_1 = \begin{bmatrix} 0.50 & 0.70 & 0.00 \\ 0.00 & 0.90 & 0.00 \\ 0.00 & 0.80 & 0.40 \end{bmatrix}, \quad \Lambda_2 = \begin{bmatrix} 0.50 & 0.67 & -0.01 \\ 0.00 & 0.94 & 0.02 \\ 0.00 & 0.00 & 0.38 \end{bmatrix}$$

$$\Rightarrow \Sigma = \begin{bmatrix} 1.33 & 0.85 & 0.00 \\ 0.85 & 22.85 & 0.60 \\ 0.00 & 0.60 & 1.19 \end{bmatrix},$$

$$\Rightarrow \Sigma \in \mathbf{M}_1 \cap \mathbf{M}_2.$$

Generic identifiability

- $\mathbf{M}_G$ admits a well-defined **algebraic dimension**.

Generic identifiability

$$\dim(\mathbf{M}_1 \cap \mathbf{M}_2) < \max\{\dim(\mathbf{M}_1), \dim(\mathbf{M}_2)\}.$$

Interpretation: $\mathbf{M}_1 \cap \mathbf{M}_2$ is a Lebesgue measure zero subset of $\mathbf{M}_1$ or $\mathbf{M}_2$.

Direct conclusions:

- Different dimensions $\Rightarrow$ identifiable.
- Knowing the dimension $\Rightarrow$ focusing on identifiability of models with the same dimension.

Jacobian matrix and Model dimension

The parametrization map:

$$\phi_G : (\Lambda, \omega) \mapsto \Sigma, \text{ s.t. } \Sigma = \Lambda^T \Sigma \Lambda + \omega I_n.$$

The Jacobian matrix of the model is

$$\mathbf{J}_G = \left(\frac{\partial (\phi_G)_j}{\partial \theta_i} \right), \quad 1 \leq i \leq E_G + 1, \quad 1 \leq j \leq \frac{n(1+n)}{2},$$

where E_G is the number of edges in G , $(\phi_G)_j$'s are the distinct entries of Σ , and θ_i 's are the entries of Λ and ω .

Proposition

$$\dim(\mathbf{M}_G) = \text{rank}(\mathbf{J}_G) \text{ at a generic point}^{(*)}$$

Proof: It's sufficient to prove that ϕ_G is a rational map.

⇒ A tool to calculate model dimension.

Linear matroid

Linear matroid of a matrix A

The pair $\{E, \mathcal{I}\}$, where $E = \{1, \dots, n\}$, representing the n columns of A , and

$$\mathcal{I} = \{S \subseteq E \mid A^S \text{ are linearly independent}\}.$$

Example: Let

$$A = \begin{bmatrix} 2 & 0 & 2 & 4 \\ 0 & 1 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ 4 & 0 & 4 & 8 \end{bmatrix},$$

then the linear matroid of A is $\{E, \mathcal{I}\}$, where

- $E = \{1, 2, 3, 4\}$;
- $\mathcal{I} = \{\emptyset, \{1\}, \{2\}, \{3\}, \{4\}, \{1, 2\}, \{1, 3\}, \{2, 3\}, \{2, 4\}, \{3, 4\}, \{1, 2, 3\}, \{2, 3, 4\}\}.$

Jacobian matroid and identifiability criteria

Jacobian matroid $\mathcal{J}(\mathbf{M}_G)$ = the **linear matroid of the Jacobian matrix**.

Proposition¹

Assume that $\dim(\mathbf{M}_1) \geq \dim(\mathbf{M}_2)$. If there exists a subset S of the coordinates such that

$$S \in \mathcal{J}(\mathbf{M}_2) \setminus \mathcal{J}(\mathbf{M}_1),$$

then $\dim(\mathbf{M}_1 \cap \mathbf{M}_2) < \min\{\dim(\mathbf{M}_1), \dim(\mathbf{M}_2)\}$.

A direct result: if $\dim(\mathbf{M}_1) = \dim(\mathbf{M}_2)$, then the condition can be transformed to

$$S \in \mathcal{J}(\mathbf{M}_1) \setminus \mathcal{J}(\mathbf{M}_2) \text{ or } S \in \mathcal{J}(\mathbf{M}_2) \setminus \mathcal{J}(\mathbf{M}_1).$$

1. Sullivant (2023)

Main results

Maximal class

Definition

A maximal class of a directed graph is the set of reachable nodes from a Strongly Connected Component (SCC) with in-degree zero.

Link to algorithm: https://github.com/Bi-xuan/maximal_class


Example:

- SCCs:

$\{1, 2, 3\}, \{4\}, \{5\}, \{6\};$

- Sources: $\{1, 2, 3\}$ and $\{5\}$ (SCCs with in-degree zero);
- Set of *maximal classes* (unique):

$\{\{1, 2, 3, 4, 6\}, \{4, 5, 6\}\}.$



Jacobian matrix structure

The Jacobian matrix $\mathbf{J}_G$ is of size $(E_G + 1) \times \frac{n(n+1)}{2}$, where E_G is the number of edges in G .

Indexing:

- **Rows:** Non-zero elements in Λ and the parameter ω .
- **Columns:** Distinct values of the covariance matrix Σ .

Proposition: Maximal class and Jacobian nullity

A column of the Jacobian matrix $\mathbf{J}_G^{[\sigma_{ij}]}$ is zero if and only if nodes i and j belong to distinct maximal classes.

Model dimension

Theorem 1: model dimension

If G does not contain multi-edges, then

$$\dim(\mathbf{M}_G) = \min \{n_r, n'_c\},$$

where

$$n_r = E_G + 1;$$

$$n'_c = |\{\{a, b\} \mid a, b \in [n], a, b \text{ belong to the same maximal class}\}|.$$

Idea of proof: We prove that the submatrix $\mathbf{J}'_G$ containing all non-zero columns of the Jacobian matrix is full rank generically by showing

1. **Non-triviality:** Prove that $\det(\mathbf{J}'_G)$ is a non-trivial polynomial.
2. **Existence:** Identify a specific parameter point where the determinant is non-zero.

Maximal class characterization

Theorem 2: Models with the same dimension

$\mathbf{M}_1$ and $\mathbf{M}_2$ with the same dimension are generically identifiable if G_1 and G_2 have different maximal classes.

Idea of proof: For models of the same dimension, generic identifiability is established by showing

$$\exists S_1 \in \mathcal{J}(\mathbf{M}_1) \setminus \mathcal{J}(\mathbf{M}_2) \text{ or } \exists S_2 \in \mathcal{J}(\mathbf{M}_2) \setminus \mathcal{J}(\mathbf{M}_1).$$

The result follows by a consequence of the main proposition ensuring that $\exists i, j \in [n]$ s.t.

$$\mathbf{J}_1^{[\sigma_{ij}]} = \mathbf{0} \text{ and } \mathbf{J}_2^{[\sigma_{ij}]} \neq \mathbf{0}.$$

Maximal class characterization

Theorem 3: Models with unknown dimensions

$\mathbf{M}_1$ and $\mathbf{M}_2$ are generically identifiable if both

1. there exist $i, j \in [n]$ s.t. i, j belong to the same maximal class in G_1 , but do not in G_2 , and
2. there exist $s, t \in [n]$ s.t. s, t belong to the same maximal class in G_2 , but do not in G_1 .

Idea of proof: We prove the following stronger criterion of Jacobian matroids difference, and the rest of the proof follows the same technique as Theorem 2.

$$\exists S_1 \in \mathcal{J}(\mathbf{M}_1) \setminus \mathcal{J}(\mathbf{M}_2) \text{ and } \exists S_2 \in \mathcal{J}(\mathbf{M}_2) \setminus \mathcal{J}(\mathbf{M}_1).$$


Summary

$\mathbf{M}_1$ and $\mathbf{M}_2$ are generically identifiable if

Condition		Example	
G_1 and G_2 do not contain multi-edges	$\dim(\mathbf{M}_1) = \dim(\mathbf{M}_2)$, $\mathcal{M}\mathcal{C}_1 \neq \mathcal{M}\mathcal{C}_2$	G_1	G_2
	$\dim(\mathbf{M}_1) \neq \dim(\mathbf{M}_2)$	G_1	G_2
G_1 or G_2 contains multi-edges	Theorem 3 is satisfied	G_1	G_2

Summary

$\mathbf{M}_1$ and $\mathbf{M}_2$ do not satisfy the sufficient identifiability criteria in this paper if

Condition		Example	
G_1 and G_2 do not contain multi-edges	$\dim(\mathbf{M}_1) = \dim(\mathbf{M}_2)$ and $\mathfrak{Mc}_1 = \mathfrak{Mc}_2$	G_1 	G_2 
G_1 or G_2 contains multi-edges	Theorem 3 is not satisfied	G_1 	G_2 is any graph

Takeaway message

1. **Novelty in identifiability context:** Focus on the stationary setting rather than time series.
2. **Rigorous and extendable algebraic framework.**
3. **Practical identifiability criteria:** Connect identifiability to graph structure, yielding straightforward criteria.
4. **Relevance to ecological research:** Incorporates the sampling insufficiency problem, requiring only i.i.d. steady-state samples instead of costly time series.

Ongoing work

Optimization

Assumption: Identifiability criteria are satisfied.

Goal: Support recovery of Λ

Recover the support of Λ and ω from Σ by minimizing the objective function:

$$f_{\Sigma}(\Lambda) = \|\Sigma - \Lambda^T \Sigma \Lambda - \omega I_n\|_F.$$

Ground truth: The true (Λ^*, ω^*) satisfies $f_{\Sigma}(\Lambda^*, \omega^*) = 0$.

Algorithms investigated:

1. **ISTA:** Fast, but low accuracy even in identifiable cases.
2. **Improved greedy algorithm:** Slow, but 100% accuracy in identifiable cases.

Stats

Assumption: Identifiability criteria are satisfied.

Plug the empirical covariance matrix $\hat{\Sigma}$ into the objective function:

$$\gamma_n(\Lambda, \omega) = \|\hat{\Sigma} - \Lambda^T \hat{\Sigma} \Lambda - \omega I\|_F^2.$$

The estimator $(\hat{\Lambda}, \hat{\omega}) = \arg \min_{(\Lambda, \omega)} \gamma_n(\Lambda, \omega)$ is **consistent**.

Sparsity-inducing estimation

In practice, we minimize the following to encourage sparsity in Λ :

$$\gamma_n(\Lambda, \omega) + \text{pen}_n(m)$$

where $\text{pen}_n(m)$ penalizes the **cardinality** of the support of Λ .

Penalization design: Based on model selection method introduced in [1, 2, 3].

1. Birgé et al. (2001)
2. Laurent et al. (2000)
3. Lebarbier (2005)

References i



I. Akjouj, M. Barbier, M. Clenet, W. Hachem, M. Maïda, F. Massol, J. Najim, and V. C. Tran.

Complex systems in ecology: a guided tour with large Lotka–Volterra models and random matrices.

Proceedings of the Royal Society A, 480(2285):20230284, 2024.



P. Legendre, M. De Cáceres, and D. Borcard.

Community surveys through space and time: testing the space–time interaction in the absence of replication.

Ecology, 91(1):262–272, 2010.



S. Chaffron, E. Delage, M. Budinich, D. Vintache, N. Henry, C. Nef, M. Ardyna, A. A. Zayed, P. C. Junger, P. E. Galand, et al.

Environmental vulnerability of the global ocean epipelagic plankton community interactome.

Science Advances, 7(35):eabg1921, 2021.

References ii



M. Gong, K. Zhang, B. Schoelkopf, D. Tao, and P. Geiger.
Discovering temporal causal relations from subsampled data.
International Conference on Machine Learning, 1898–1906, 2015.



A. Tank, E. B. Fox, and A. Shojaie.
Identifiability and estimation of structural vector autoregressive models for subsampled and mixed-frequency time series.
Biometrika, 106(2):433–452, 2019.



W. C. Young, K. Y. Yeung, and A. E. Raftery.
Identifying dynamical time series model parameters from equilibrium samples, with application to gene regulatory networks.
Statistical modelling, 19(4):444–465, 2019.



M. Drton, B. Hollering, and J. Wu.
Identifiability of homoscedastic linear structural equation models using algebraic matroids.
Advances in Applied Mathematics, 163:102794, 2025.

References iii



L. Birgé and P. Massart.

Gaussian model selection.

Journal of the European Mathematical Society, 3(3):203–268, 2001.



B. Laurent and P. Massart.

Adaptive estimation of a quadratic functional by model selection.

Annals of statistics, 1302–1338, 2000.



É. Lebarbier.

Detecting multiple change-points in the mean of Gaussian process by model selection.

Signal processing, 85(4):717–736, 2005.

Thank you for your attention!